



# Tabla de dosis para RCP en perros y gatos



		Peso (kg)	2.5	5	10	15	20	25	30	35	40	45	50
DROGA		DOSIS	ml	ml	ml	ml	ml	ml	ml	ml	ml	ml	ml
Paro	Epinefrina/adrenalina (1:1000; 1mg/mL)	0,01 mg/kg	0.03	0.05	0.1	0.15	0.2	0.25	0.3	0.35	0.4	0.45	0.5
	Vasopresina (20 U/mL)	0,8 U/kg	0.1	0.2	0.4	0.6	0.8	1	1.2	1.4	1.6	1.8	2
	Atropina (0,4 - 0,54 mg/mL)	~ 0,05 mg/kg	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
Anti-Arrítmico	Amiodarona (50 mg/mL)	5 mg/kg	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	Lidocaína (20 mg/mL)	2 mg/kg	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	Esmolol* (10 mg/mL)	0,5 mg/kg	0.13	0.25	0.5	0.75	1	1.3	1.5	1.8	2	2.3	2.5
Reversión	Naloxona (0,4 mg/mL)	0,04 mg/kg	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	Flumazenil (0,1 mg/mL)	0,01mg/kg	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	Atipamezol (5 mg/mL)	100 µg/kg	0.06	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
Desfibrilación monofásica	Desfibrilador externo (J)	4 - 6 J/kg	10 J	20 J	40 J	60 J	80 J	100 J	120 J	140 J	160 J	180 J	200 J
	Desfibrilador interno (J)	0,5 - 1 J/kg	2 J	3 J	5 J	8 J	10 J	15 J	15 J	20 J	20 J	20 J	25 J
		Peso (kg)	2.5	5	10	15	20	25	30	35	40	45	50

\*Administrar esmolol 0,5 mg/kg IV o IO durante 3-5 minutos seguido de una IRC a 50 mcg/kg/min.

# Tabla de dosis para RCP en perros y gatos



		Peso (kg)	2.5	5	10	15	20	25	30	35	40	45	50
<b>DROGA</b>		<b>DOSIS</b>	ml	ml	ml	ml	ml	ml	ml	ml	ml	ml	ml
<b>Paro</b>	<b>Epinefrina/adrenalina</b> (1:1000; 1mg/mL)	<b>0,01 mg/kg</b>	0.03	0.05	0.1	0.15	0.2	0.25	0.3	0.35	0.4	0.45	0.5
	<b>Vasopresina</b> (20 U/mL)	<b>0,8 U/kg</b>	0.1	0.2	0.4	0.6	0.8	1	1.2	1.4	1.6	1.8	2
	<b>Atropina</b> (0,4 - 0,54 mg/mL)	<b>~ 0,05 mg/kg</b>	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
<b>Anti-Arrítmico</b>	<b>Amiodarona</b> (50 mg/mL)	<b>5 mg/kg</b>	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	<b>Lidocaína</b> (20 mg/mL)	<b>2 mg/kg</b>	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	<b>Esmolol*</b> (10 mg/mL)	<b>0,5 mg/kg</b>	0.13	0.25	0.5	0.75	1	1.3	1.5	1.8	2	2.3	2.5
<b>Reversión</b>	<b>Naloxona</b> (0,4 mg/mL)	<b>0,04 mg/kg</b>	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	<b>Flumazenil</b> (0,1 mg/mL)	<b>0,01mg/kg</b>	0.25	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	<b>Atipamezol</b> (5 mg/mL)	<b>100 µg/kg</b>	0.06	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
<b>Desfibrilación bifásica</b>	<b>Desfibrilador externo</b> (J)	<b>2 - 4 J/kg</b>	5 J	10 J	20 J	30 J	40 J	50 J	60 J	70 J	80 J	90 J	100 J
	<b>Desfibrilador interno</b> (J)	<b>0,2 - 0,4 J/kg</b>	1 J	2 J	2 J	3 J	4 J	5 J	6 J	7 J	8 J	9 J	10 J
		Peso (kg)	2.5	5	10	15	20	25	30	35	40	45	50

\*Administrar esmolol 0,5 mg/kg IV o IO durante 3-5 minutos seguido de una IRC a 50 mcg/kg/min.